

A Heavy Higgs and a Light Sneutrino NLSP in the MSSM with Enhanced $SU(2)$ D-terms

Anibal D. Medina ^a, Nausheen R. Shah ^{b,d}
and Carlos E. M. Wagner ^{b,c,d}

Department of Physics, University of California, One Shields Ave. Davis, CA 95616 ^a, USA

Enrico Fermi Institute ^b and Kavli Institute for Cosmological Physics ^c,

University of Chicago, , 5640 S. Ellis Ave., Chicago, IL 60637, USA

HEP Division, Argonne National Laboratory, 9700 Cass Ave., Argonne, IL 60439, USA ^d

April 10, 2009

Abstract

The Minimal Supersymmetric extension of the Standard Model provides a solution to the hierarchy problem and leads to the presence of a light Higgs. A Higgs boson with mass above the present experimental bound may only be obtained for relatively heavy third generation squarks, requiring a precise, somewhat unnatural balance between different contributions to the effective Higgs mass parameter. It was recently noticed that somewhat heavier Higgs bosons, which are naturally beyond the LEP bound, may be obtained by enhanced weak $SU(2)$ D-terms. Such contributions appear in models with an enhanced electroweak gauge symmetry, provided the supersymmetry breaking masses associated with the scalars responsible for the breakdown of the enhanced gauge symmetry group to the Standard Model one, are larger than the enhanced symmetry breaking scale. In this article we emphasize that the enhanced $SU(2)$ D-terms will not only raise the Higgs boson mass but also affect the spectrum of the non-standard Higgs bosons, sleptons and squarks, which therefore provide a natural contribution to the T parameter, compensating for the negative one coming from the heavy Higgs boson. The sleptons and non-standard Higgs bosons of these models, in particular, may act in a way similar to the so-called inert Higgs doublet. The phenomenological properties of these models are emphasized and possible cosmological implications as well as collider signatures are described.

1 Introduction

The Minimal Supersymmetric Standard Model (MSSM) provides an elegant solution to the hierarchy problem of the Standard Model (SM) and in the presence of R-parity, naturally includes a good dark matter candidate. The MSSM also leads to the presence of a Higgs boson, with SM-like properties, which is naturally light, with a tree-level mass smaller than the Z -gauge boson mass [1, 2]. Therefore, the current experimental bound on a SM-like Higgs boson from the LEP experiment, $m_h \gtrsim 114.4$ GeV, demands relatively large radiative corrections induced by the third generation squarks [3],[4]. Such large corrections may only be obtained by large squark masses which lead at the same time to large negative radiative corrections to the soft supersymmetry breaking Higgs mass parameter. Hence, proper electroweak symmetry breaking (EWSB) may only be obtained by fine-tuning the negative soft supersymmetry breaking and positive μ term contributions to the effective Higgs mass parameter.

The situation may be improved if the electroweak SM gauge group proceeds from a larger gauge structure which is spontaneously broken to the SM group at energies of a few TeV. If the supersymmetry breaking parameter of the scalars, which are associated with the breakdown of the extended gauge symmetry, is much larger than the order parameter of the gauge symmetry breakdown, the D-term of the low energy weak interactions may be enhanced. This, in turn, leads to larger values of the tree-level SM-like Higgs boson mass and reduces fine tuning [5],[6],[7].

A natural implementation of these ideas may be realized by extending the weak gauge group to the group $SU(2)_1 \times SU(2)_2$, where the two Higgs doublets of the MSSM as well as the third generation particles are charged under one of the two $SU(2)$ groups, $SU(2)_1$, while the first and second generation particles are charged under $SU(2)_2$ [8]. The $SU(2)_1 \times SU(2)_2$ symmetry is eventually broken to the diagonal $SU(2)_W$ group that may be identified with the one associated with the SM weak interactions. Such an extension of the MSSM is anomaly free and can lead to an understanding of the hierarchy of the third generation and the lighter generation masses. For relatively large values of the $SU(2)_1$ gauge coupling one can get SM-like Higgs boson masses up to 300 GeV.

Such large Higgs boson masses tend to be in conflict with precision electroweak observables [9]. The model described above, provides extra contributions to these observables. On one side, there are non-universal contributions to the third generation quark and lepton couplings to the weak gauge bosons, which are induced by the mixing of the heavy gauge bosons with the SM ones. Such contributions may be minimized for large values of the vacuum expectation value (vev) of the scalars that provide the breakdown of the extended symmetry (see, for example, Ref. [10]). However, the presence of a heavy Higgs still leads to a relatively large negative contribution to the T -parameter, leading to further tension with the precision electroweak observables.

In this work we emphasize that the same D-term contributions that lead to an enhancement of the Higgs mass also lead to a splitting of the left-handed super-partners of the third generation squarks and sleptons as well as the non-standard Higgs bosons. We show that the mass splitting of these doublets is naturally of the size necessary to compensate the large negative Higgs contribution, restoring agreement of the model with precision electroweak data, similar to the inert Higgs doublet models [12]. The modification of the spectrum leads to novel phenomenological properties with

respect to the MSSM. It is the aim of this article to discuss these properties as well as the associated supersymmetry breaking mechanism that may be consistent with such a spectrum.

The article is organized as follows. In section 2 we review the model. In section 3 we describe the Higgs and sparticle spectrum and discuss the contributions to the T -parameter. In section 4 we describe a possible realization of supersymmetry breaking that leads to a spectrum of third generation left-handed sparticles much lighter than the first and second generation ones and discuss the collider physics associated with this model. In section 5 we comment on the possible cosmological constraints. In section 6 we describe a different possibility, in which the non-standard Higgs bosons become light, while the left-handed squarks and sleptons remain heavy. We reserve section 7 for our conclusions.

2 Review of the Model

A simple way to implement the possibility of an enhanced electroweak D-term is through the enlarged weak gauge group $SU(2)_1 \times SU(2)_2$. The two MSSM Higgs bosons and the third generation left-handed doublet super-fields are charged under $SU(2)_1$ but singlets under $SU(2)_2$, while the lighter generations left-handed doublets are singlets under $SU(2)_1$ but charged under $SU(2)_2$. A bidoublet, Σ , is introduced which acquires a vev, $\langle \Sigma \rangle = uI$, breaking the product group to the diagonal one, $SU(2)_W$, which is identified with the SM weak group. The effective weak gauge coupling is therefore given by

$$g = \frac{g_1 g_2}{\sqrt{g_1^2 + g_2^2}}, \quad (1)$$

where g_1 and g_2 are the $SU(2)_1$ and $SU(2)_2$ gauge couplings, respectively. For values of $g_1 \gg g_2$ the weak coupling is approximately equal to g_2 . Following Ref [5], the breakdown of the enhanced gauge symmetry is governed by the Σ superpotential

$$W = \lambda_1 S \left(\frac{\Sigma \Sigma}{2} - w^2 \right), \quad (2)$$

where S is a gauge singlet. This leads to a Σ potential of the form

$$V = m_\Sigma^2 \Sigma^\dagger \Sigma + \frac{\lambda_1^2}{4} |\Sigma \Sigma|^2 - \frac{B}{2} (\Sigma \Sigma + h.c.) + \dots \quad (3)$$

where $B = \lambda_1 w^2$, m_Σ^2 is a soft supersymmetry breaking mass term and, for simplicity, we have considered all parameters to be real. There is also a contribution coming from the D-terms of the $SU(2)_1$ and $SU(2)_2$ gauge groups to the potential,

$$\Delta V = \frac{g_1^2}{8} \left(\text{Tr}[\Sigma^\dagger \tau^a \Sigma] + H_u^\dagger \tau^a H_u + H_d^\dagger \tau^a H_d + L^\dagger \tau^a L + Q^\dagger \tau^a Q \right)^2 + \frac{g_2^2}{8} \left(\text{Tr}[\Sigma^\dagger \tau^a \Sigma] + \dots \right)^2 \quad (4)$$

where τ^a are the generators of $SU(2)$ and, for completeness, we have added the extra third generation lepton and quark scalars, which were omitted in the analysis of Ref. [5].

For values of $B > m_\Sigma^2$, the Σ field acquires a vev along the D-flat direction, $\langle \Sigma \rangle = uI$, with $u^2 = (B - m_\Sigma^2)/\lambda_1^2$. Assuming that $B \gg v^2$, with v the vev of the SM Higgs boson, one can integrate out the heavy degrees of freedom and find the result for the enhanced $SU(2)_W$ D-terms

$$\Delta V = \frac{g^2}{2} \Delta \sum_a \left(H_u^\dagger \tau^a H_u + H_d^\dagger \tau^a H_d + L_3^\dagger \tau^a L_3 + Q_3^\dagger \tau^a Q_3 \right)^2 \quad (5)$$

and

$$\Delta = \frac{1 + \frac{2m_\Sigma^2}{g_2^2 u^2}}{1 + \frac{2m_\Sigma^2}{(g_2^2 + g_1^2) u^2}}. \quad (6)$$

Therefore, ignoring mixing with the non-standard CP-even Higgs boson, the SM-like Higgs mass is enhanced to a value

$$m_h^2 = \frac{1}{2} (g^2 \Delta + g_Y^2) v^2 \cos^2 2\beta + \lambda_2^2 v^2 \sin^2 2\beta + \text{loop corrections} \quad (7)$$

which is the main result of Ref. [5]. For completeness, we added the possible contribution of a superpotential term

$$W_\lambda = \lambda_2 S' H_u H_d \quad (8)$$

with S' a singlet superfield. In our analysis we shall assume that the D-terms give the dominant contribution to the tree-level Higgs mass. The term λ_2 , however, may have important phenomenological implications as have been observed in several works, including Refs. [6, 11]. In our phenomenological analysis, we will consider the simple case in which $\lambda_2 = 0$.

Assuming that the value of u is of the order of a few TeV, as required to minimize the non-oblique corrections to precision electroweak observables, the authors of Ref. [5] showed that the presence of the enhanced D-terms can naturally raise the Higgs mass up to values of about 250 GeV, and for large values of the g_1 coupling at the symmetry breaking scale, $\alpha_1 \simeq 1$, values as large as 300 GeV may be obtained.

3 Sparticle Spectrum and Phenomenological Implications

An important aspect of this model that was omitted from Ref. [5] is that the presence of the D-terms will also induce additional splitting in the Higgs, squark and slepton masses. In order to understand this point, one can write the low energy $SU(2)$ D-term contributions to the effective potential in the following way

$$V_D = \frac{g^2 \Delta}{8} \left(\sum_i \Phi_i^\dagger \Phi_i \right)^2 - \frac{g^2 \Delta}{4} \sum_{ij} |\Phi_i^T i \sigma_2 \Phi_j|^2, \quad (9)$$

where $\Phi_i = L_3, Q_3, H_i$ are the $SU(2)$ doublets with non-trivial transformations under $SU(2)_1$ and σ_2 is the antisymmetric Pauli matrix. To obtain the spectrum one should concentrate on

interactions of the different particles with the Higgs fields. The first term in Eq. (9) leads to a positive contribution, $g^2\Delta v^2/4$ to the mass of the third generation squarks and sleptons, where $v^2 = v_u^2 + v_d^2$, and $v_{u,d}$ are the $H_{u,d}$ vevs. The second term in V_D is the sum of the squares of the $SU(2)$ invariant combinations that couple to the corresponding quark and lepton $SU(2)$ singlets in the superpotential Yukawa terms. Therefore, it leads to negative contributions, $-g^2\Delta v_i^2/2$, to the squark and slepton doublet component that couples via the Yukawa interaction with the Higgs H_i , with $i = u, d$.

The sparticles also receives F -term contributions associated with the superpotential

$$W = \mu H_u H_d + h_u H_u Q U + h_d H_d Q D + h_l H_d L E. \quad (10)$$

In particular, the third generation squarks receive supersymmetric contributions proportional to their super-partner masses, which become particularly important in the case of the top squarks. The combination of the F-term and D-term contributions lead to the following splitting between the third generation left-handed squark and slepton mass parameters,

$$m_{\tilde{\tau}_L}^2 - m_{\tilde{\nu}_\tau}^2 = \Delta_D \quad (11)$$

$$m_{\tilde{b}_L}^2 - m_{\tilde{t}_L}^2 = \Delta_D - m_t^2 \quad (12)$$

and

$$\begin{aligned} \Delta_D &= \frac{g^2 v^2}{2} \Delta |\cos 2\beta| \\ &= (\Delta m_h^2)_D / |\cos 2\beta| \end{aligned} \quad (13)$$

where $v \simeq 174.1$ GeV is the SM Higgs boson vev and $(\Delta m_h^2)_D = g^2 v^2 \Delta \cos^2(2\beta)/2$ is the enhanced D-term contribution to the tree-level Higgs mass and we have ignored all SM quark and lepton masses apart from the top quark mass. For moderate to large values of $\tan \beta$, $\tan \beta > 5$, which we will assume from now on, $\Delta_D \simeq (\Delta m_h^2)_D$.

The case of the non-standard Higgs bosons should be treated separately. The charged Higgs and CP-odd states are defined as orthogonal to the corresponding charged and neutral Goldstone boson states, namely

$$\begin{aligned} H^\pm &= \sin \beta H_d^\pm - \cos \beta H_u^\pm \\ A &= \sin \beta A_d - \cos \beta A_u \end{aligned} \quad (14)$$

where $H_i^\pm$ and A_i are the charged Higgs and CP-odd components of the Higgs field doublets H_i ($i = u, d$), and for H_u we redefine $H_u \rightarrow i\sigma_2 H_u^*$ which carries the same hypercharge as H_d .

Similar to the MSSM case, the neutral CP-odd mass is given by

$$m_A^2 = m_1^2 + m_2^2 \quad (15)$$

where $m_1^2 = m_{H_d}^2 + |\mu|^2$ and $m_2^2 = m_{H_u}^2 + |\mu|^2$, where $m_{H_d}^2$ and $m_{H_u}^2$ are the soft supersymmetry breaking square mass parameters. The resultant mass splitting is given by

$$m_{H^\pm}^2 - m_A^2 = \frac{g^2 \Delta}{2} v^2. \quad (16)$$

3.1 Contributions to the T -Parameter

The additional splitting described in the previous section between the masses of the Higgs bosons, sleptons and squarks can have important phenomenological implications. For instance, contrary to naive expectations, for large values of the D-term contribution to the Higgs mass, the left-handed sbottom becomes heavier than the left-handed stops. The exact stop and sbottom spectrum depends, however, on the mixing with the right-handed third generation squarks. In this section, to simplify the discussion, we will assume those mixings to be small, something that is natural for non-degenerate squarks, moderate values of A_t and non-extreme values of $\tan\beta$.

The splitting between the upper and lower components of the left-handed doublets control their contribution to the T parameter. For an $SU(2)$ doublet, with up and down mass eigenvalues m_u and m_d , the contribution to the T -parameter is given by

$$\Delta T = \frac{N_c}{16\pi s_W^2 m_W^2} \left[m_u^2 + m_d^2 - \frac{2m_u^2 m_d^2}{m_u^2 - m_d^2} \log \left(\frac{m_u^2}{m_d^2} \right) \right], \quad (17)$$

where s_W is the sine of the standard weak mixing angle $s_W^2 \simeq 0.2315$.

If the ratio of the heavier doublet mass eigenstate to the lighter one is smaller than about 3, the contribution to the T parameter may be approximated by [12]

$$\begin{aligned} \Delta T &= \frac{N_c}{12\pi s_W^2 m_W^2} (\Delta m_{ud})^2 \\ &= \frac{N_c}{12\pi s_W^2 m_W^2} \frac{(\Delta m_{ud}^2)^2}{(m_u + m_d)^2}, \end{aligned} \quad (18)$$

where N_c is the number of colors, Δm_{ud} and Δm_{ud}^2 are the differences between the masses and the squared masses of the heavier and lighter components of the doublet. These contributions should be added to the ones associated with the heavy SM-like Higgs boson,

$$\begin{aligned} \Delta T &= -\frac{3}{8\pi c_W^2} \ln \frac{m_h}{m_{h_{\text{ref}}}} \\ \Delta S &= \frac{1}{6\pi} \ln \frac{m_h}{m_{h_{\text{ref}}}}, \end{aligned} \quad (19)$$

where $m_{h_{\text{ref}}}$ is a Higgs mass reference value. As mentioned above, the up and down sfermion mass eigenstates are admixtures of $SU(2)$ doublet and singlet components. In the case of non-negligible mixing, the expression for ΔT , given by Eq. (18), needs to be reformulated. This turns out to be specially important for third generation sfermions. In the case of an $SU(2)$ doublet, taking into account mixing angles, the general expression for ΔT takes the form,

$$\begin{aligned} \Delta T &= \frac{N_c}{12\pi s_W^2 m_W^2} \left(\sin^2 \theta_u \sin^2 \theta_d (m_{u_2} - m_{d_2})^2 \right. \\ &\quad + \sin^2 \theta_u \cos^2 \theta_d (m_{u_2} - m_{d_1})^2 + \cos^2 \theta_u \sin^2 \theta_d (m_{u_1} - m_{d_2})^2 \\ &\quad + \cos^2 \theta_u \cos^2 \theta_d (m_{u_1} - m_{d_1})^2 - \sin^2 \theta_u \cos^2 \theta_u (m_{u_2} - m_{u_1})^2 \\ &\quad \left. - \sin^2 \theta_d \cos^2 \theta_d (m_{d_2} - m_{d_1})^2 \right), \end{aligned} \quad (20)$$

where θ_u and θ_d are the mixing angles for the up and down component of the $SU(2)$ doublet respectively, and $m_{u_{1,2}}$, $m_{d_{1,2}}$ are the mass eigenvalues. In the case when all masses are of the same order or we have zero mixing, the approximation used in Eq. (18) may be applied.




Figure 1: Values of ΔT necessary to provide a good fit to electroweak precision data, for two values of the Higgs mass: $m_h = 150$ GeV and $m_h = 300$ GeV.

We present in Fig. 1 the necessary values of new physics contributions to the S and T parameters, to get agreement with precision electroweak observables at the 1σ level. The values used here¹ include the recent measurements of $m_W = 80.432 \pm 0.039$ GeV and $m_t = 172.4 \pm 1.2$ GeV, and are consistent with the ones obtained in Ref. [9]. The precise values of S and T , however, depend on the observables used in the fit (see for example Ref. [13]). From now on, we will use the values displayed in Fig. 1 as our reference values.

The scalars give only a small contribution to the S -parameter, hence the ellipses provide us with information about the values of ΔT . As can be seen from Fig. 1, for Higgs boson masses in the range $m_h \simeq 150$ – 300 GeV, the necessary value of ΔT to provide a good fit to the precision electroweak observables is

$$\begin{aligned}\Delta T_{150} &\simeq 0.10 \pm 0.06 \\ \Delta T_{300} &\simeq 0.24 \pm 0.07\end{aligned}\tag{21}$$

¹We are thankful to Jens Erler for providing us with the fit used to generate these values.

where ΔT_{m_h} is the extra contribution to the T parameter necessary to provide a good fit to the precision electroweak data with a Higgs of mass m_h (GeV).

Before considering the implications for the sparticles and Higgs masses one should stress that the enhanced D-term contribution is obtained from a small deviation of the Σ scalar from the $SU(2)_W$ singlet u -direction. The triplet vev leads therefore to a contribution to the T parameter

$$\Delta T \simeq \frac{4\pi g_1^4}{s_W^2 c_W^2 g^4} \frac{m_W^2 u^2}{M_T^4} \quad (22)$$

where M_T is the triplet mass. For values of u and the triplet mass necessary to significantly enhance the tree-level Higgs mass, these contributions prove to be too small to compensate for the negative Higgs mass contributions to the T -parameter for $m_h \lesssim 300$ GeV. Furthermore, it is clear from the numbers presented in Eq. (21) and Fig. 1, that for SM-like Higgs masses smaller than 150 GeV, the need for new physics contributions to the precision measurements, and therefore the constraints on the sparticle spectrum, become weaker. From now on, we will hence concentrate on the region $m_h \gtrsim 150$ GeV, although somewhat smaller Higgs masses are still consistent with the presence of enhanced weak $SU(2)$ D-terms.

If sufficiently light, the third generation squark and sleptons, as well as the non-standard Higgs bosons can lead to large contributions to the T parameter. The squark contribution is enhanced by a color factor, but is suppressed by the cancellation between the D-term and F-term contributions to the left-handed sbottom-stop splitting. Moreover, in the simplest soft supersymmetry breaking schemes, the squarks become much heavier than the sleptons and, from Eq. (18) their contributions to the T -parameter are further suppressed. We shall first assume that the non-standard Higgs boson mass is a few times larger than the third generation left-handed slepton masses, something that is realized, for instance, in the simple scenario to be discussed in the following section. This means that the non-standard Higgs bosons will lead to small contributions to the T -parameter compared to the slepton one.

Considering that only the sleptons contribute to the T parameter, the requirement of restoring agreement with precision electroweak observables defines the allowed range of slepton masses. Using the values obtained in Fig. 1 for a Higgs mass of 150 GeV, ignoring mixing in the squark and slepton sectors and assuming that the Higgs mass receives two-loop contributions from third generation squarks with masses of about ~ 1 TeV, this leads, for the lower (upper) required values of the new physics contribution to the T parameter, to sneutrino and stau masses:

$$\begin{aligned} m_{\tilde{\nu}} &\simeq 150 \text{ (30) GeV} \\ m_{\tilde{\tau}} &\simeq 195 \text{ (120) GeV.} \end{aligned} \quad (23)$$

For a Higgs mass of 300 GeV, instead, we get a relatively heavier spectrum

$$\begin{aligned} m_{\tilde{\nu}} &\simeq 380 \text{ (260) GeV} \\ m_{\tilde{\tau}} &\simeq 480 \text{ (395) GeV.} \end{aligned} \quad (24)$$

The sleptons should then acquire masses of the order of a few hundred GeV, with a relatively large mass splitting, which grows for larger Higgs masses and is in the range of a few tens of GeV to more

than a 100 GeV. The results are shown in Figs. 2, 3 and 4. In Fig. 3 we have also drawn a straight line with slope of $m_h/2$. Sneutrinos below the line are potential candidates for Higgs decays.




Figure 2: Range of preferred values of the stau mass [GeV] as a function of the Higgs mass [GeV].

Furthermore, there exists a lower bound on the sneutrino mass, $m_{\tilde{\nu}_\tau} \gtrsim 40$ GeV, coming from the invisible decay width of the Z gauge boson measured at LEP[13]. This bound, as shown in Fig. 3, is almost automatically fulfilled for sneutrino masses in the range selected by precision electroweak measurements for $m_h > 150$ GeV included in the plots.

4 Phenomenological Properties

4.1 Higgs Boson Searches

One of the most important differences of this model with the standard implementation of the MSSM is the Higgs mass range, which has an important impact on its searches. For the mass range $150 \text{ GeV} \lesssim m_h \lesssim 300 \text{ GeV}$ that we are considering in this article, the most important SM-like Higgs boson search channels will be its decays into charged and neutral gauge bosons, namely

$$\begin{aligned}
 H &\rightarrow W^\pm W^\mp \\
 H &\rightarrow Z Z.
 \end{aligned} \tag{25}$$




Figure 3: Range of preferred values of the tau-sneutrino mass [GeV] as a function of the Higgs mass [GeV]. Sneutrinos below the black line have masses less than $m_h/2$ and therefore are potential candidates for Higgs decays.

These searches will become very efficient in the first years of the LHC run [14], and therefore we expect these kind of models to be probed with the first few fb^{-1} of LHC running. Moreover, if the Higgs mass is below 190 GeV, then even the Tevatron will be able to probe these models in the coming years [15]. Indeed, the Tevatron has already excluded the existence of a SM-like Higgs boson with mass between 160 and 170 GeV at the 95% confidence level, and is expected to probe the whole range of masses, $m_h \in [150, 190]$ GeV, by the end of next year.

The Higgs boson searches would change if there would be new, supersymmetric decays of the Higgs bosons. If we assume that the lightest superpartner of the SM particles is the light tau-sneutrino discussed in the last section, there is only a small region of parameters where these decays would be open, and only for masses below 190 GeV, that coincides with the range to be explored by the Tevatron collider. As can be seen from Fig. 3, for larger values of the Higgs mass, the tau-sneutrino is sufficiently heavy as to avoid any Higgs decay into light supersymmetric particles.

For Higgs boson masses below 190 GeV the sneutrinos may be light enough to allow on-shell decays of the Higgs into two sneutrinos. Moreover, the coupling of third generation sneutrinos to the Higgs boson is enhanced with respect to the one in the MSSM, due to the enhanced $SU(2)$




Figure 4: Range of preferred values of the stau-sneutrino mass difference [GeV] as a function of the Higgs mass [GeV].

D-term contribution and is given by

$$g_h \tilde{\nu}_\tau \tilde{\nu}_\tau \simeq -i \frac{(g^2 \Delta + g_Y^2) v}{2\sqrt{2}}. \quad (26)$$

Due to this enhanced coupling, the Higgs boson may have a significant decay branching ratio into light sneutrinos, therefore avoiding the Tevatron bounds. More quantitatively, assuming masses above the $2 m_W$ threshold, the decay width for decays into two vector bosons is given by

$$\Gamma(h \rightarrow VV) \simeq \frac{G_F(|Q_V| + 1) m_h^3}{\sqrt{2} 16 \pi} \left(1 - \frac{4m_V^2}{m_h^2} + \frac{12m_V^4}{m_h^4}\right) \left(1 - \frac{4m_V^2}{m_h^2}\right)^{1/2} \quad (27)$$

where G_F is the Fermi constant, Q_V is the charge of the massive gauge boson, $V = W^\pm$, Z , and M_V is its mass. For the decay into sneutrinos, instead, the result is

$$\Gamma(h \rightarrow \tilde{\nu}_\tau \tilde{\nu}_\tau) \simeq \frac{(g^2 \Delta + g_Y^2)^2 v^2}{128 \pi m_h} \left(1 - \frac{4m_{\tilde{\nu}_\tau}^2}{m_h^2}\right)^{1/2}. \quad (28)$$

For instance, for a Higgs mass $m_h \simeq 165$ GeV and a sneutrino mass of about 70 GeV, the branching ratio of the Higgs into a pair of $W^\pm$ gauge bosons is reduced to less than half of its SM

value, avoiding the current Tevatron bound. Somewhat smaller or larger values of the sneutrino masses, within the range consistent with precision measurements and smaller than a half of the Higgs mass, lead to more or less restrictive Tevatron bounds on this model, respectively.

4.2 Supersymmetric Particle Spectrum

In this section, we will work in the moderate $\tan\beta$ limit, which for universal scalar masses, tend to lead to heavy non-standard Higgs bosons and is therefore close to the so-called decoupling limit in the Higgs sector. In the foregoing discussion we assumed that the mixing of left-handed and right-handed third generation sleptons is small. Whether the lightest stau is mostly left-handed or right-handed depends on the supersymmetry breaking mechanism. We will assume that $\tilde{\tau}_1$ is mostly left-handed while $\tilde{\tau}_2$ is mostly right-handed, and present a mechanism that realizes this possibility below.

The tree-level third generation Yukawa couplings are given by

$$h_t = \frac{gm_t}{\sqrt{2}m_W \sin\beta}, \quad h_b = \frac{gm_b}{\sqrt{2}m_W \cos\beta}, \quad h_\tau = \frac{gm_\tau}{\sqrt{2}m_W \cos\beta}. \quad (29)$$

As is well known, h_b and h_τ can become comparable to h_t in the large $\tan\beta$ regime. As previously mentioned, we are interested in the moderate $\tan\beta$ limit and therefore only the top quark Yukawa will be of importance. In the examples found later in the text, we take the reference value of $\tan\beta = 10$.

We shall assume that Yukawa couplings for the first and second generation quarks and leptons are generated by adding a massive Higgs-like pair of doublets which transform under $SU(2)_2$, as was suggested in [5], and work with this minimal spectrum. With this minimal spectrum, the β_i function coefficients are given by

$$b_Y = \frac{36}{5}, \quad b_{22} = 1, \quad b_{21} = -1, \quad b_3 = -3, \quad (30)$$

where b_{22} and b_{21} correspond to the $SU(2)_2$ and $SU(2)_1$ gauge groups, while b_Y and b_3 correspond to the hypercharge and strong interactions, respectively. The above values should be compared to the MSSM case:

$$b_Y = \frac{33}{5}, \quad b_2 = 1, \quad b_3 = -3. \quad (31)$$

This will generally upset unification of couplings [16]. Let us stress, however, that for the chosen low energy values of these couplings, the $SU(2)_2$ coupling α_2 becomes close to α_Y and α_3 at scales of about the standard grand unification scale, $M_{GUT} \approx 2 \times 10^{16}$ GeV.

The modified β function coefficients also have consequences for the soft supersymmetry breaking parameters. For instance, assuming a common value for the gaugino mass, $M_{1/2}$, at the messenger scale, the running of the Bino mass is modified, making the lightest neutralino lighter than in the MSSM for the same value of $M_{1/2}$. Regarding the third generation masses, if we assume a common gaugino mass, they will be pushed to large values driven by the strong $SU(2)_1$ gaugino effects. Therefore, for the third generation sleptons to remain light, we need a mechanism of SUSY

breaking that keeps the third generation soft scalar masses small at low energies while the strong $SU(2)_1$ coupling g_1 runs to large values.

We can accomplish this with a model where SUSY breaking is transmitted to the visible sector via only the $SU(3)_c \times SU(2)_2 \times U(1)_Y$ gauginos. The $SU(2)_1$ gauge sector then does not contribute to the running of soft third generation masses. For simplicity, the input parameters for the model are: $\tan \beta$; a universal gaugino mass $M_{1/2}$ for the $SU(3)_c$, $SU(2)_2$ and $U(1)_Y$ gauginos; a universal scalar mass m_0 for all squarks and sleptons; soft-supersymmetry breaking parameters m_{H_u} and m_{H_d} for the two Higgs bosons which interact under $SU(2)_1$; and we assume positive $\text{sign}(\mu)$ and $A_t = A_b = A_\tau = 0$ at the messenger scale M which is of the order of M_{GUT} . The $SU(2)_1$ gaugino remains massless, but acquires a mass of order $g_1 u$, together with the $SU(2)_1$ gauge bosons, after the breakdown of $SU(2)_1 \times SU(2)_2 \rightarrow SU(2)_W$.

Lets take a look at the one-loop renormalization group equations (RGE) for the left- and right-handed soft stau masses at energy scales larger than the gauge symmetry breaking scale $SU(2)_1 \times SU(2)_2 \rightarrow SU(2)_W$,

$$16\pi^2 \frac{d}{dt} m_{L_3}^2 = -\frac{6}{5} g_Y^2 |M_Y|^2 - \frac{3}{5} g_Y^2 S \quad (32)$$

$$16\pi^2 \frac{d}{dt} m_{\tilde{\tau}_R}^2 = -\frac{24}{5} g_Y^2 |M_Y|^2 + \frac{6}{5} g_Y^2 S \quad (33)$$

where S is given by,

$$S \equiv \text{Tr}[Y_i m_{\phi_i}^2] = m_{H_u}^2 - m_{H_d}^2 + \text{Tr}[m_Q^2 - m_L^2 - 2m_u^2 + m_d^2 + m_e^2]. \quad (34)$$

In Eqs.(32) and (33) we have neglected the terms proportional to the τ -Yukawa coupling, which remain small unless $\tan \beta$ is very large and, as stressed above, we have assumed that our symmetry breaking scheme is $SU(2)_1$ blind. Furthermore we take $m_{H_u} = m_{H_d}$ and since we have chosen a universal scalar mass m_0 at high energy, this implies that S vanishes and does not run with energy. Therefore, given that the $U(1)_Y$ charges for left-handed sleptons are smaller than the $U(1)_Y$ charges of the right-handed sleptons, this naturally leads to the left-handed stau soft masses being smaller than the right-handed ones, as can be seen from Eqs.(32) and (33). This is in contrast to what happens in the MSSM [17, 18, 19], where the non-universal requirement $m_{H_d} - m_{H_u} > 0$ is necessary to obtain smaller left-handed masses for the third generation sleptons. In the case of soft gaugino masses, their one-loop RGE take the form,

$$16\pi^2 \frac{d}{dt} M_i^2 = 4b_i g_i^2 M_i^2 \quad (35)$$

where, as stressed above, up to the $SU(2)_1 \times SU(2)_2$ breaking scale $b_i = (36/5, -1, 1, -3)$ for the $U(1)_Y$, $SU(2)_1$, $SU(2)_2$ and $SU(3)_c$ gauge groups, and after the breakdown of $SU(2)_1 \times SU(2)_2 \rightarrow SU(2)_W$, we get the standard MSSM result, $b_i = (33/5, 1, -3)$.

The solution of the RG equations show that the soft supersymmetry breaking parameters $m_{L_3}^2$ and $m_{\tilde{\tau}_R}^2$ are approximately given by

$$m_{L_3}^2 \simeq m_0^2 + 0.04 M_{1/2}^2, \quad m_{\tilde{\tau}_R}^2 \simeq m_0^2 + 0.15 M_{1/2}^2, \quad (36)$$

while the low energy Bino and Wino masses are given by

$$M_Y \simeq 0.35 M_{1/2}, \quad M_2 \simeq 0.8 M_{1/2}^2. \quad (37)$$

The requirement of getting the appropriate stau and tau-sneutrino masses for a given value of the Higgs mass and $\tan\beta$ defines the allowed parameter space in the m_0 – $M_{1/2}$ plane. In particular, looking at Eq.(18) and (36), we notice that for a fixed Higgs mass and in the absence of stau mixing there is an ellipsoidal area in the $M_{1/2}$ – m_0 plane where our spectrum fits electroweak precision data at a 68 % confidence level. In Fig. 5 we plot this correlation for several Higgs masses neglecting the stau mixing. For large values of $|\mu|$, there is a lower bound of $M_{1/2} \gtrsim 130$ GeV coming from the LEP bound on chargino masses [20]. We see that the preferred region of parameter space is relatively thin and moves to higher soft masses as the Higgs mass increases, roughly maintaining its thickness. The red (medium light gray), blue (dark gray) and purple (medium dark gray) regions in the figure have the additional constraint of a $\tilde{\nu}_\tau$ NLSP, whereas in the green (light gray) region $m_{\tilde{\chi}_0} < m_{\tilde{\nu}_\tau}$. The blue and purple regions instead, have the constraint that the bino is heavier than the stau, $m_{\tilde{\tau}} < m_{\tilde{\chi}_0}$. Finally, in the blue region the stau dominant decay mode is into an on-shell $W^\pm$ and a tau-sneutrino; this does not happen in the purple region, where it decays off-shell to $W^\pm$ and a tau-sneutrino and also into an off-shell Bino and a τ lepton. We can see from the figure that, as expected, the region where $\tilde{\nu}_\tau$ is the NLSP becomes more prominent for smaller values of m_0 . Furthermore, for larger values of the Higgs mass, the mass difference between the stau and the tau-sneutrino is sufficiently large to allow on-shell decays into a sneutrino and a charged gauge boson in most of the parameter space. We will assume in this work, as is natural if the messenger scale is somewhat smaller than the GUT scale, that the LSP is the gravitino and, as can be seen in Fig. 5 appropriate boundary conditions at high energy can be chosen such that the NLSP is the lightest tau sneutrino, $\tilde{\nu}_\tau$. This should be compared with, for instance, the standard low scale gaugino mediated scenario, in which the right-handed staus are naturally the NLSP [21, 22, 23].

Due to our mechanism of SUSY breaking and the new big D-term contributions to third generation sleptons, there is a lower bound on the soft scalar universal mass coming from the requirement of the sneutrino to be in the right mass range to lead to agreement with precision electroweak observables. This can be seen in Fig. 5, where for example, for a Higgs mass $m_h \simeq 150$ GeV, $m_0 \gtrsim 40$ GeV for $M_{1/2} = 500$ GeV to fulfill this requirement.

In the case of a universal gaugino mass, $M_{1/2}$, at high energy, we naturally obtain that the lightest neutralino, $\tilde{\chi}_1^0$, is mostly bino-like, while the second lightest neutralino, $\tilde{\chi}_2^0$, is mostly wino-like. This implies that right-handed squarks mostly decay to $q \tilde{\chi}_1^0$. From Fig. 5, depending on the initial masses, m_0 and $M_{1/2}$, we see that we can have either $m_{\tilde{\tau}_1} < m_{\tilde{\chi}_1^0}$ or $m_{\tilde{\chi}_1^0} < m_{\tilde{\tau}_1}$ for a small scalar mass m_0 and a moderate gaugino mass $M_{1/2}$, once we take into account the D-term corrections. We also expect the left-handed first and second family squarks and sleptons to be heavier than the lightest neutralino since their masses run with $SU(3)_c$ and $SU(2)_2$ couplings respectively. With these considerations in mind, for a light sneutrino NLSP, we infer the following possible mass spectra:

$$m_{\tilde{G}} < m_{\tilde{\nu}_\tau} < m_{\tilde{\tau}_1} < m_{\tilde{\chi}_1^0} < m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^\pm} < m_{\tilde{\tau}_2}, m_{\tilde{l}_R}, m_{\tilde{l}_L} \dots \quad (38)$$

$$m_{\tilde{G}} < m_{\tilde{\nu}_\tau} < m_{\tilde{\chi}_1^0} < m_{\tilde{\tau}_1} < m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^\pm} < m_{\tilde{\tau}_2}, m_{\tilde{l}_R}, m_{\tilde{l}_L} \dots \quad (39)$$




Figure 5: Correlation of m_0 and $M_{1/2}$ for different Higgs masses and $\tan\beta = 10$. Red represents the region where $\tilde{\nu}_\tau$ is the NLSP and $m_{\tilde{\tau}_1} > m_{\tilde{\chi}_0}$, blue and purple are the regions where $\tilde{\nu}_\tau$ is the NLSP and $m_{\tilde{\chi}_0} > m_{\tilde{\tau}_1}$ and staus decay predominantly to on shell Ws and tau sneutrinos (blue region) or to off-shell Ws and sneutrinos or off-shell binos and taus (purple region). Green is the region where χ_0 is the NLSP.

which are realized in Tables 1, 2 and 3.

In order to calculate the particle spectrum at low energies for some sample points at high energies and to calculate the main decays we used SDECAY [24]. SDECAY works mainly within the MSSM and thus we had to slightly modify the RGE used in the program to make it suitable for our model. From the RG running we expect modifications to the spectrum at high energies. Furthermore SDECAY only calculates two body decays of sparticles. To understand the behavior at low energies, let us write the tree level stau mass matrix, taking into account the non-zero trilinear terms and the left-right mixing,

$$m_{\tilde{\tau}}^2 = \begin{pmatrix} m_{L_3}^2 + \Delta_{L_3} - \frac{g^2}{4}(\Delta - 1)v^2 \cos 2\beta & m_{\tau}(A_{\tau} - \mu \tan \beta) \\ m_{\tau}(A_{\tau} - \mu \tan \beta) & m_{\tilde{\tau}_R}^2 + \Delta_{e_3} \end{pmatrix} \quad (40)$$

where m_{L_3} and $m_{\tilde{\tau}_R}$ are the third generation left- and right-handed slepton soft breaking masses respectively, $\Delta_{L_3} = (-1/2 + \sin^2 \theta_w) \cos 2\beta m_Z^2$ and $\Delta_{e_3} = \sin^2 \theta_w \cos 2\beta m_Z^2$. We can see that the mass eigenstates will have an admixture of left- and right-handed components, mainly due to the mixing caused by the μ term in Eq.(40). However, for $\tan \beta \simeq 10$, due to the smallness of the τ mass, we expect this admixture to be small. Similarly, the tau sneutrino modified mass is given by,

$$m_{\tilde{\nu}_{\tau}}^2 = m_{L_3}^2 + \Delta_{\nu_{\tau}} + \frac{g^2}{4}(\Delta - 1)v^2 \cos 2\beta \quad (41)$$

where $\Delta_{\nu_{\tau}} = \cos 2\beta m_Z^2/2$.

Furthermore, minimizing the tree level Higgs potential we find the following expression for μ ,

$$\mu^2 = \frac{1}{2} \left(\frac{m_{H_u}^2 - m_{H_d}^2}{\cos 2\beta} - m_{H_u}^2 - m_{H_d}^2 - m_{\Delta}^2 \right) \quad (42)$$

where $m_{\Delta}^2 = (g_1^2 + g_2^2 \Delta)v^2/2$, and likewise the tree-level Higgs masses are given by,

$$m_A^2 = \frac{m_{H_u}^2 - m_{H_d}^2}{\cos 2\beta} - m_{\Delta}^2, \quad (43)$$

$$m_{h^0}^2 = \frac{1}{2} \left(m_A^2 + m_{\Delta}^2 - \sqrt{(m_A^2 - m_{\Delta}^2)^2 + 4m_{\Delta}^2 m_A^2 \sin 2\beta} \right), \quad (44)$$

$$m_{H^0}^2 = \frac{1}{2} \left(m_A^2 + m_{\Delta}^2 + \sqrt{(m_A^2 - m_{\Delta}^2)^2 + 4m_{\Delta}^2 m_A^2 \sin 2\beta} \right), \quad (45)$$

$$m_{H^+}^2 = m_A^2 + \frac{g^2}{2} \Delta v^2. \quad (46)$$

Observe that for $\Delta = 1$, the above expressions reduce to the well known MSSM ones.

4.3 Collider Signatures

The particle spectrum and the most important branching ratios for some sample points are presented in Tables 1, 2 and 3. Table 1 presents a case of sneutrinos which are light enough to allow the decay

of a Higgs into two on-shell tau-sneutrinos. Similar to the example discussed in the previous section, the Higgs decay branching ratio into a pair of $W^\pm$'s is reduced to a value that is less than half of the SM result, therefore avoiding the current Tevatron bounds.

As can be seen from Table 2 and 3, two body decays, $\tilde{\tau}_1^\pm \rightarrow W^\pm \tilde{\nu}_\tau$, tend to be kinematically allowed and thus may be an important decay modes for the light staus. Therefore, we expect relatively hard leptons and/or jets plus missing energy to be the an important signature from $\tilde{\tau}_1$ decays. This is very different from regular gaugino mediated SUSY breaking scenarios in the MSSM where the lightest stau tends to decay into τ 's and gravitinos, and the heaviest stau, since the mass difference, $m_{\tilde{\tau}} - m_{\tilde{\nu}_\tau}$ is small, tends to decay into an on-shell τ and a Bino. Indeed, the small sneutrino-stau mass difference allows only 3-body decays, which through an off-shell W , leads to soft fermions: $f\bar{f}'$ plus missing energy [18].

Sparticle	Mass[GeV]	Dominant decay modes
$\tilde{g}$	1146	$\tilde{q}_L q$ (18)%, $\tilde{q}_R q$ (36)%, $\tilde{b}_{1,2} b$ (23) %, $\tilde{t}_1 t$ (23) %
$\tilde{u}_L, \tilde{d}_L$	1047, 1050	$\tilde{\chi}_2^0 q$ (32) %, $\tilde{\chi}_1^\pm q'$ (64) %
$\tilde{u}_R, \tilde{d}_R$	1010, 1007	$\tilde{\chi}_1^0 q$ (99) %
$\tilde{t}_1$	804	$\tilde{\chi}_1^+ b$ (34) %, $\tilde{\chi}_1^0 t$ (27) %, $\tilde{\chi}_2^0 t$ (14) %
H^+	724	
A	708	
$\tilde{\chi}_4^0$	644	$\tilde{\chi}_1^\pm W^\mp$ (54) %, $\tilde{\chi}_2^0 h$ (17) %
$\tilde{\chi}_2^\pm$	644	$\tilde{\chi}_2^0 W^\pm$ (28) %, $\tilde{\chi}_1^\pm Z$ (26) %, $\tilde{\chi}_1^\pm h$ (18) %
$\tilde{\chi}_3^0$	630	$\tilde{\chi}_1^\pm W^\mp$ (57) %, $\tilde{\chi}_2^0 Z$ (25) %
$\tilde{\chi}_2^0$	385	$\tilde{\nu}_\tau \nu_\tau$ (48) %, $\tilde{\tau}_1^\pm \tau^\mp$ (41) %
$\tilde{\chi}_1^\pm$	385	$\tilde{\nu}_\tau \tau^\pm$ (50) %, $\tilde{\tau}_1^\pm \nu_\tau$ (38) %
$\tilde{e}_L$	346	$\tilde{\chi}_1^0 e$ (100) %
$\tilde{\nu}_e$	337	$\tilde{\chi}_1^0 \nu_e$ (100) %
$\tilde{e}_R$	209	$\tilde{\chi}_1^0 e$ (100) %
$\tilde{\tau}_2$	218	$\tilde{\chi}_1^0 \tau$ (72) %, $\tilde{\nu}_\tau W$ (27) %
$\tilde{\chi}_1^0$	178	$\tilde{\nu}_\tau \nu_\tau$ (83) %, $\tilde{\tau}^\pm \tau^\mp$ (17)%
$\tilde{\tau}_1$	140	
$\tilde{\nu}_\tau$	73	$\tilde{G} \nu_\tau$

Table 1: Spectrum and branching ratios for $M_{1/2} = 500$ GeV, $\tan \beta = 10$, $m_{H_u}^2 = m_{H_d}^2 = 10^4 \text{GeV}^2$, $m_0 = 90$ GeV, $\Delta = 3.57$ and $m_h = 169$ GeV. Since the first and second generations are almost degenerate, we only give results for the first and third sfermion generations.

Among the major differences with the case of non-universal Higgs masses (NUHM) [17] [18], [19] is that the masses of the first two generation families run with $SU(2)_2$ making left-handed sleptons and squarks heavier than right handed ones, whereas in the NUHM case, for all generations, since S is positive, it tends to make left-handed sleptons of all families relatively light. This is important since it affects the decay of the charginos and the second lightest neutralinos, which, contrary to the

Sparticle	Mass[GeV]	Dominant decay modes
$\tilde{g}$	1564	$\tilde{q}_L q$ (16.2)%, $\tilde{q}_R q$ (31.4)%, $\tilde{b}_{1,2} b$ (20) %, $\tilde{t}_1 t$ (24) %
$\tilde{u}_L, \tilde{d}_L$	1428, 1429	$\tilde{\chi}_2^0 q$ (32) %, $\tilde{\chi}_1^\pm q'$ (64) %
$\tilde{u}_R, \tilde{d}_R$	1374, 1368	$\tilde{\chi}_1^0 q$ (99) %
$\tilde{t}_1$	1112	$\tilde{\chi}_1^+ b$ (19) %, $\tilde{\chi}_1^0 t$ (25) %, $\tilde{\chi}_3^0 t$ (17) %, $\tilde{\chi}_2^+ b$ (23) %
H^+	967	
A	946	
$\tilde{\chi}_4^0$	864	$\tilde{\chi}_1^\pm W^\mp$ (56) %, $\tilde{\chi}_2^0 h$ (19) %
$\tilde{\chi}_2^\pm$	864	$\tilde{\chi}_2^0 W^\pm$ (28) %, $\tilde{\chi}_1^\pm Z$ (28) %, $\tilde{\chi}_1^\pm h$ (20) %
$\tilde{\chi}_3^0$	852	$\tilde{\chi}_1^\pm W^\mp$ (56) %, $\tilde{\chi}_2^0 Z$ (26) %
$\tilde{\chi}_2^0$	551	$\tilde{\nu}_\tau \nu_\tau$ (47) %, $\tilde{\tau}_1^\pm \tau^\mp$ (39) %
$\tilde{\chi}_1^\pm$	551	$\tilde{\nu}_\tau \tau^\pm$ (49) %, $\tilde{\tau}_1^\pm \nu_\tau$ (37) %
$\tilde{e}_L$	486	$\tilde{\chi}_1^0 e$ (100) %
$\tilde{\nu}_e$	480	$\tilde{\chi}_1^0 \nu_e$ (100) %
$\tilde{e}_R$	300	$\tilde{\chi}_1^0 e$ (100) %
$\tilde{\tau}_2$	303	$\tilde{\chi}_1^0 \tau$ (72) %, $\tilde{\nu}_\tau W$ (28) %
$\tilde{\chi}_1^0$	249	$\tilde{\nu}_\tau \nu_\tau$ (90) %, $\tilde{\tau}_1^\pm \tau^\mp$ (10) %
$\tilde{\tau}_1$	217	$\tilde{\nu}_\tau W$ (100) %
$\tilde{\nu}_\tau$	132	$\tilde{G} \nu_\tau$

Table 2: Spectrum and branching ratios for $M_{1/2} = 700$ GeV, $\tan \beta = 10$, $m_{H_u}^2 = m_{H_d}^2 = 10^4 \text{GeV}^2$, $m_0 = 150$ GeV, $\Delta = 6.13$ and $m_h = 210$ GeV. Since the first and second generations are almost degenerate, we only give results for the first and third sfermion generations.

Sparticle	Mass[GeV]	Dominant decay modes
$\tilde{g}$	829	$\tilde{q}_R q$ (42)%, $\tilde{b}_{1,2} b$ (16) %, $\tilde{t}_1 t$ (42) %
$\tilde{u}_L$	853	$\tilde{\chi}_2^+ q'$ (46) %, $\tilde{\chi}_4^0$ (23) %, $\tilde{\chi}_1^+ q'$ (15) %
$\tilde{d}_L$	857	$\tilde{\chi}_2^- q'$ (51) %, $\tilde{\chi}_4^0$ (25) %
$\tilde{u}_R, \tilde{d}_R$	750, 737	$\tilde{\chi}_1^0$ (92) %
H^+	620	
A	588	
$\tilde{t}_1$	539	$\tilde{\chi}_1^+ b$ (79) %, $\tilde{\chi}_1^0 t$ (20) %
$\tilde{\chi}_4^0$	580	$\tilde{\chi}_1^\pm W^\mp$ (27) %, $\tilde{\tau}_1^\pm \tau^\mp$ (22) %, $\tilde{\nu}_\tau \nu_\tau$ (30) %
$\tilde{\chi}_2^\pm$	580	$\tilde{\chi}_2^0 W^\pm$ (13) %, $\tilde{\chi}_1^\pm Z$ (28) %, $\tilde{\chi}_3^0 W^\pm$ (10) %, $\tilde{\nu}_\tau \tau^\pm$ (28) %, $\tilde{\tau}_1^\pm \nu_\tau$ (24) %
$\tilde{\chi}_3^0$	400	$\tilde{\chi}_1^0 Z$ (88) %
$\tilde{\chi}_2^0$	387	$\tilde{\tau}_2^\pm \tau^\mp$ (14) %, $\tilde{\tau}_1^\pm \tau^\mp$ (54) %
$\tilde{\chi}_1^\pm$	378	$\tilde{\nu}_\tau \tau^\pm$ (41) %, $\tilde{\chi}_1^0 W^\pm$ (47) %
$\tilde{e}_L$	495	$\tilde{\chi}_1^0 e$ (65) %, $\tilde{\chi}_2^0 e$ (20) %, $\tilde{\chi}_2^0 e$ (20) %, $\tilde{\chi}_1^\pm \nu_e$ (14) %
$\tilde{\nu}_e$	489	$\tilde{\chi}_1^0 \nu_e$ (80) %, $\tilde{\chi}_1^\pm e^\mp$ (18) %
$\tilde{e}_R$	302	$\tilde{\chi}_1^0 e$ (100) %
$\tilde{\tau}_2$	299	$\tilde{\chi}_1^0 \tau$ (94) %
$\tilde{\chi}_1^0$	249	$\tilde{\nu}_\tau \nu_\tau$ (99) %
$\tilde{\tau}_1$	242	$\tilde{\nu}_\tau W$ (100) %
$\tilde{\nu}_\tau$	163	$\tilde{G} \nu_\tau$

Table 3: Spectrum and branching ratios for $M_Y = M_2 = 700$ GeV and $M_3 = 350$ GeV at the messenger scale, $\tan \beta = 10$, $m_{H_u}^2 = m_{H_d}^2 = 1 \times 10^4 \text{GeV}^2$, $m_0 = 150$ GeV, $\Delta = 6.13$ and $m_h = 208$ GeV. Since the first and second generations are almost degenerate, we only give results for the first and third sfermion generations.




Figure 6: Examples of typical squark decays.

NUHM, now decay predominantly into only third generation sleptons. The consequence is events with τ leptons in the final state, without a counterpart of similar events with electrons or muons.

In the case of first and second generation squarks, seen in Tables 1 and 2, as happens in the MSSM, the right-handed squarks are still the lightest ones. This leads, in the case of heavy gluinos, to smaller branching ratio of gluinos into left-handed squarks than right-handed ones. This, in turn, implies a bigger production of binos which tend to decay more prominently into the sneutrino NLSP. Since the bino decay products are invisible particles, the corresponding collider signatures resemble those of the standard bino LSP scenarios. This is partially compensated by the lighter third generation squarks, which tend to decay into charginos and second lightest neutralinos, which, in turn decay into third generation sleptons. Moreover, as seen in Fig. 5, there are large regions of parameter space where $m_{\tilde{\chi}_0^1} > m_{\tilde{\tau}_1}$, giving a non-negligible decay branching ratio of binos into lighter staus which later decay into either on-shell or off-shell W 's. This is particularly true for larger values of the Higgs mass. Examples of typical decay cascades expected at the LHC or the Tevatron are given in Fig. 6. For these, we find, using Table 2, that the total branching ratios are 10 %, 12 % and 31%, respectively. In the case that the decays start with a gluino, after considering the extra jet, the latter branching ratios become 3.14×10^{-2} , 1.94×10^{-2} and 5×10^{-2} , respectively.

The existence of a heavy Higgs can lead to very interesting events, with many $W^\pm$'s and bottom quarks in the final state. This feature can be seen in Table 2 where a potential channel for Higgs production in supersymmetric particle decays arises. This channel is given by the decay chain $\tilde{g} \rightarrow \tilde{t}_1 t \rightarrow \tilde{\chi}_2^+ b t \rightarrow \tilde{\chi}_1^+ h b t$. In the last stage of the decay chain we can have $\tilde{\chi}_1^+$ decaying either directly to the NLSP, $\tilde{\nu}_\tau \tau$, about half the time or to $\tilde{\tau}_1 \nu_\tau$ about 37 % of the time, the latter case being more phenomenologically interesting since $\tilde{\tau}_1 \rightarrow W \tilde{\nu}_\tau$ produces on-shell $W^\pm$'s. Therefore, we can have the following final states $\tilde{g} \rightarrow h b W b \tilde{\nu}_\tau \tau$ and $\tilde{g} \rightarrow h b W b \nu_\tau W \tilde{\nu}_\tau$ with gluino decay branching ratios of 5.4×10^{-3} or 4.3×10^{-3} , respectively. Each gluino therefore may decay into a maximum of 2 bottom quarks and 4 $W^\pm$ states, plus missing energy. A pair of $W^\pm$ should reconstruct the Higgs mass, while the invariant mass of one of the bottom quarks and one of the $W^\pm$'s should be equal to the top quark mass.

One can also consider the possibility of searching for the heavy non-standard Higgs bosons, via their decay into SM-like Higgs bosons. This was analyzed in Ref. [25] in a similar supersymmetric model, in which the heavy Higgs boson masses are induced by a large coupling to a singlet field [6],[11]. This search may be possible, provided the non-standard Higgs decays into bottom and tau pairs and/or the sneutrino decay branching ratio of standard and non-standard Higgs bosons does not become too-large, and is therefore more efficient for smaller values of $\tan \beta$ and/or larger sneutrino masses.

Table 3 represents the case of a more compressed spectrum, obtained by reducing the gluino mass with respect to the Bino and Wino masses at the messenger scale [26]. This leads to the presence of relatively light stops, radically changing the phenomenology, since now a large fraction of the gluinos can decay into light stops and right-handed squarks and negligibly into left-handed squarks. In addition, the lightest stau becomes heavier than the second lightest neutralino and lightest chargino, which are Higgsino-like and couple strongly to the third generation squarks. This has the consequence of making most decays end in either $\tilde{\chi}_1^0$ or $\tilde{\nu}_\tau$ without passing through $\tilde{\tau}_1$. Observe that when searching for stops at the LHC, the main decays of $\tilde{t}_1$ in this sample point are

70% into a bottom quarks and a W gauge boson (including 20% into an on-shell top quarks) and missing energy and 30% into a tau lepton, a bottom quark and missing energy.

If we want to be in the region of parameter space where the tau sneutrino is the NLSP, from Fig. 5 and Tables 1, 2 and 3, we see that the weak gauginos usually tend to be heavier than sleptons in the low energy theory. Therefore in the case of universal gaugino masses, we predict that the first and second generation sleptons will mainly decay into the lightest neutralino, that may be important for the associated direct production phenomenology at either lepton or hadron colliders. Observe that in Table 3 due to the smallness of M_3 , the μ parameter is suppressed compared to the universal gaugino case. Therefore, as mentioned above, the lightest chargino as well as the second and third lightest neutralino are mainly Higgsino-like.

We would like to conclude this subsection by considering the effects of sfermion mixing in electroweak precision data. One of the consequences of mixing is to lower the mass of the lightest eigenstate with respect to the un-mixed state. Therefore, the analysis presented in Fig. 5 of the regions of parameter space of consistent ΔT , slightly underestimates the region where the lightest $\tilde{\tau}$ is lighter than the Bino. In the specific examples analyzed using SDECAY, given in Tables 1, 2 and 3, we confirmed that the mixing does not change the general results obtained in the case of zero mixing. We calculated the contributions to ΔT coming from the mass splitting in staus, third generation squarks and the charged Higgs sector. These contributions are shown in Table 4, where

m_h [GeV]	m_0 [GeV]	$M_{1/2}$ [GeV]	$\Delta T_{\tilde{\tau}}$	$\Delta T_{\tilde{Q}_3}$	$\Delta T_{H^\pm}$	ΔT_{tot}
169	90	500	1.5×10^{-1}	8.7×10^{-4}	7.8×10^{-4}	1.56×10^{-1}
210	150	700	1.9×10^{-1}	2.6×10^{-3}	8×10^{-3}	2.03×10^{-1}
210	150	(700,350)	1.5×10^{-1}	2.4×10^{-2}	1.9×10^{-2}	1.98×10^{-1}

Table 4: Complete ΔT calculation for particular points in parameter space, which lie in the 68 % confidence level ellipse in ΔS – ΔT plane.

we see that the results lie in the 68 % confidence level ellipses in the ΔS – ΔT plane discussed in Fig. 1, and where in the third entry for the third sample point we mean $M_1 = M_2 = 700$ GeV and $M_3 = 350$ GeV at the messenger scale. Observe that in this last case, the third generation squarks give a relatively large contribution to the T parameter.

5 Cosmological Constraints

Within the supersymmetry breaking scheme described in the last section, the tau-sneutrino is naturally the lightest SM superpartner. However, as stressed before, if the messenger scale is smaller than the standard GUT scale, the lightest supersymmetric particle tends to be the gravitino. There could be potentially constraints to this scenario specifically from the observed Dark Matter (DM) density and from the decays of the sneutrino or bino NLSP. First of all let us mention that in the case of a long lived neutralino NLSP, obtained for a large messenger scale of the order of M_{GUT} , the constraints coming from its late decay into a photon and a gravitino are strong enough to exclude

all parameter space [27]. However, if the lightest neutralino is the LSP and not the gravitino, this scenario could survive and give the correct DM relic density via co-annihilation of the LSP with a light sneutrino as discussed in [29].

In the case of the sneutrino NLSP, the dominant decay channel is,

$$\tilde{\nu}_\tau \rightarrow \tilde{G} + \nu_\tau \quad (47)$$

with a decay rate given by,

$$\Gamma_{\tilde{\nu}_\tau \rightarrow \tilde{G} + \nu_\tau} = \frac{m_{\tilde{\nu}_\tau}^5}{48\pi M_P^2 m_{\tilde{G}}^2} \left(1 - \frac{m_{\tilde{G}}^2}{m_{\tilde{\nu}_\tau}^2}\right)^4. \quad (48)$$

From this expression we can estimate the lifetime of the NLSP as a function of the sneutrino and the gravitino masses. If the gravitino mass is in the region $[1, 100]$ GeV, as it is if the messenger scale is of the order of the standard GUT scale, the lifetime of the NLSP makes it potentially dangerous for BBN for NLSP masses $\sim [45, 400]$ GeV, the latter range being the sneutrino masses of interest. Dependence of the sneutrino lifetime on the gravitino mass can be seen in Fig. 7. Assuming a




Figure 7: Lifetime of $\tilde{\nu}_\tau$ (sec) as a function of $m_{\tilde{\nu}_\tau}$ (GeV). The blue and red curves correspond to $m_{\tilde{G}} = 40$ GeV and $m_{\tilde{G}} = 1$ GeV, respectively

gravitino LSP, its relic density is related to the sneutrino NLSP density prior to decay by,

$$\Omega_{\tilde{G}} h^2 = \frac{m_{\tilde{G}}}{m_{\tilde{\nu}_\tau}} \Omega_{\tilde{\nu}_\tau} h^2 + \Omega_{\tilde{G}}^T h^2. \quad (49)$$

Given a sneutrino mass and assuming thermal equilibrium, its freeze-out temperature is roughly $T \sim m_{\tilde{\nu}_\tau}/25$. The relevant annihilation channels have already been calculated in detail in the

literature, (see for instance, Ref. [17]), the most important being Z -mediated s-channel annihilation to fermions. After a simple calculation, one obtains, for a tau sneutrino mass of about a few hundred GeV, a thermal relic density, $\Omega_{\tilde{\nu}_\tau} h^2 \approx \mathcal{O}(10^{-3})$.

Since we are interested in the case $m_{\tilde{G}} < m_{\tilde{\nu}_\tau}$, we see from Eq. (49) and from the sneutrino relic density that in order to obtain a gravitino DM relic density in accordance with experiments, we must rely on either thermal or non-thermal production of gravitinos at the reheating epoch. Gravitinos are produced in the primordial plasma right after reheating from scattering processes of the type, $x + y \rightarrow \psi + z$ where x , y and z are relevant particles and ψ is the gravitino. The longitudinal component of the gravitino is related to the goldstino, whose interaction with other particles is proportional to $(m_{\tilde{G}} M_P)^{-1}$. Hence, for masses $m_{\tilde{G}} \gtrsim 100$ keV, gravitinos are not thermalized with the rest of the plasma.

Assuming scattering processes to be the main source of gravitino production, we can estimate, for a given gravitino mass, an upper bound on the reheating temperature so that gravitinos are the main component of DM. This was done in Ref. [30], where solving the Boltzmann Equation using a thermally averaged cross-section for scattering mainly off of gluinos and squarks, it was found that,

$$\Omega_{\tilde{G}} \sim 0.2 \times \left(\frac{m_{\tilde{G}}}{\text{GeV}} \right)^{-1} \times \left(\frac{M_3}{10^3 \text{GeV}} \right)^2 \times \left(\frac{T_{reh}}{10^8 \text{GeV}} \right). \quad (50)$$

Taking into account that the gluinos are usually heavy, in Fig. 8 we plotted the bound on the reheating temperature as a function of the gravitino mass, assuming that the latter saturates the DM component of the energy density in the Universe.




Figure 8: Reheating temperature, T_{reh} , as a function of the gravitino mass, $m_{\tilde{G}}$. The red and blue curves correspond to a soft gluino mass $M_3 = 500, 1000$ GeV respectively.

If the sneutrino decays during or after BBN, it could alter the predictions for the light-element abundances. This is due mainly to two effects. On one hand, if the mass difference between the sneutrino and gravitino is sufficiently large, the sneutrino decay produces high energy non-thermal neutrinos which, through scattering processes with background particles, transfer part of their energy to them. Some of the processes to consider are $\nu_\tau + \bar{\nu}_{i,BG} \rightarrow (e^\pm, \mu^\pm \tau^\pm)$, $\nu_\tau + \bar{\nu}_{i,BG} \rightarrow \pi^+ + \pi^-$. On the other hand, subdominant multi-body (bigger than two) sneutrino decays into charged and/or strongly interacting particles can also contribute significantly, despite having small branching ratios. Hadronic constraints, coming from processes such as,

$$\tilde{\nu}_\tau \rightarrow \tilde{G}\nu_\tau q\bar{q}, \quad \tilde{\nu}_\tau \rightarrow \tilde{G}\nu_\tau Z, \quad \tilde{\nu}_\tau \rightarrow \tilde{G}\tau W, \quad (51)$$

with W and Z decaying into jets, turn out to be much stronger than electromagnetic constraints, coming from processes such as,

$$\tilde{\nu}_\tau \rightarrow \tilde{G}\nu_\tau l\bar{l}, \quad \tilde{\nu}_\tau \rightarrow \tilde{G}\nu_\tau Z, \quad \tilde{\nu}_\tau \rightarrow \tilde{G}\tau W, \quad (52)$$

with W and Z decaying leptonically. The most stringent constraint comes from the overproduction of D and ${}^6\text{Li}$ produced in hadron showers induced by four-body decays [31].

There is also a potential constraint coming from the Cosmic Microwave Background (CMB). For early decays, EM cascades are thermalized completely by Compton scattering, double Compton scattering and bremsstrahlung and no spectral distortion of the CMB takes place. However for late decays, the spectrum cannot relax to thermodynamic equilibrium, resulting in a Bose-Einstein distribution with a non-vanishing chemical potential, μ_{pot} . The current constraint from experiments is $\mu_{pot} < 9 \times 10^{-5}$. This constraint turns out to be negligible in the case of a sneutrino NLSP with a gravitino LSP, as was shown in [32].

All of these effects have been studied in Refs. [31] and [32] where, assuming a thermal relic density for the sneutrino NLSP, it was found that for a gravitino with a mass $m_{\tilde{G}} \gtrsim 10$ GeV, $\Delta m = m_{\tilde{\nu}_\tau} - m_{\tilde{G}} \lesssim 290$ GeV and for $m_{\tilde{G}} \lesssim 10$ GeV, there is no constraint on Δm . As mentioned above, the biggest constraint comes from hadron showers affecting D and ${}^6\text{Li}$ BBN predictions. Therefore, since in our model $m_{\tilde{\nu}_\tau} \lesssim 300$ GeV (380 GeV) for $m_h < 270$ GeV ($m_h < 300$ GeV) (see Fig. 3) and we work with gravitino masses $1 \text{ GeV} < m_{\tilde{G}} < 100$ GeV, we conclude that it is only for the larger Higgs masses that our parameter space gets slightly truncated by cosmological constraints. Therefore most of our parameter space of interest safely passes all cosmological constraints.

6 Model with Non-vanishing $SU(2)_1$ Gaugino Masses

Throughout this article, we have assumed that the $SU(2)_1$ gaugino masses vanish at the messenger scale and therefore left handed third generation soft SUSY breaking slepton and squark masses as well as soft SUSY Higgs masses do not run under the gauge group $SU(2)_1$. Hence, the corresponding particles are not driven to large values via the $SU(2)_1$ gaugino corrections. In particular third generation sleptons remain light as a consequence of the transmission of SUSY breaking mechanism via only $SU(3)_c \times SU(2)_2 \times U(1)_Y$ gaugino masses.

We also saw how the doublet spectrum is split by $SU(2)_1$ D-terms in the effective low energy theory. This provides a compensating contribution to the T -parameter that cancels the logarithmic contribution coming from a large Higgs mass. Which particle provides the cancellation to the standard Higgs contribution to ΔT is a model dependent question that has important consequences on the supersymmetric particle spectrum. In most of this work, we have assumed third generation sleptons to be light with respect to the third generation squarks and non-standard Higgs doublet, and therefore to also provide the main contribution to this cancellation. That the lightest doublet under $SU(2)_1$ is the main contributor to ΔT is easy to understand from looking at Eq.(18), the squared mass difference is independent of the soft masses and thus ΔT is inversely proportional to the sparticle masses.

If we drop our assumption that the third generation sleptons, squarks and Higgs soft masses do not run under $SU(2)_1$, then their masses are pushed in an analogous fashion to larger values at low energies by the strong g_1 coupling. For both third generation squarks and sleptons there are small Yukawa effects that lower their respective masses, but do not compensate the strong $SU(2)_1$ and $SU(3)_c$ gaugino contributions. Therefore, we conclude that in the case of $SU(3)_c \times SU(2)_1 \times SU(2)_2 \times U_Y$ gaugino SUSY breaking mediation, it will be mostly up to the non-standard Higgs doublet to provide the main contribution to ΔT . Although we will not make a detailed phenomenological analysis, we will present the most important phenomenological properties.

Let us analyze this case in slightly more depth. According to Eq. (43), the CP-odd Higgs mass depends only on the difference of the soft supersymmetry breaking parameters of the Higgs doublets $m_{H_{u,d}}^2$. Therefore, for universal soft scalar masses, the CP-odd and charged Higgs become light in the large $\tan\beta$ limit [33], when the top and bottom Yukawa couplings affecting the Higgs mass parameters become of the same order. Moreover, since now we have the contributions from $SU(2)_1$ gauginos to the sparticle RGEs, right-handed sfermions will tend to be lighter than left-handed ones. In particular, due to the enhancement of the third generation slepton and Higgs mass parameters, the right-handed stau will be affected by large tau-Yukawa effects, pushing it to small values. Thus, the phenomenology will be similar to the light stau NLSP in gaugino mediation models. An example of the low energy spectrum arising from this kind of models is presented in Table 5. Note that large values of $\tan\beta$ and the $SU(2)_1$ gaugino mass, M_1 , may lead to dangerously small values of the right-handed stau mass due to the negative Yukawa corrections induced by the heavy left-handed sleptons. This is the main reason why the value of M_1 in the example of Table 5 was chosen to be smaller than the rest of the gaugino masses.

Contrary to what happens in the case of vanishing $SU(2)_1$ gaugino masses, the light CP-odd scalars demanded in this scenario are restricted by direct searches at the Tevatron collider and by flavor constraints (see for example Ref. [34]). For instance, for values of $\tan\beta \simeq 50$, a light CP-odd Higgs with a mass below 170 GeV is strongly constrained by current Tevatron bounds [35]. This restricts an important region of the parameter space corresponding to $m_h < 180$ GeV as can be observed from Fig. 3 by substituting the sneutrino mass by the CP-odd Higgs mass. All Tevatron bounds may be avoided for SM-like Higgs boson masses above 190 GeV, for which heavier CP-odd Higgs bosons may still give a relevant ΔT contribution. Future searches at the Tevatron will probe the existence of CP-odd Higgs bosons with masses up to 250 GeV for this range of values of $\tan\beta$,

and will lead to further tests of this scenario. Observe also that the flavor constraints coming from the rare decays, $B_u \rightarrow \tau \nu$ and $b \rightarrow s \gamma$, will be modified with respect to the MSSM case, due to the larger mass difference between the charged and CP-odd Higgs bosons. An analysis of this question, although very interesting, is beyond the scope of the present article.

Cosmological constraints on this scenario arise in the case of a stau NLSP with a gravitino LSP dark matter. These are related to BBN constraints in the case of gravitino masses bigger than 100 KeV and to over-closure of the Universe and/or large scale structure formation for smaller masses². For messenger scales of the order of M_{GUT} , however, the gravitino can be heavy enough so that the neutralino may be the lightest supersymmetric particle, leading to the same cosmological and phenomenological properties as the MSSM in the stau–neutralino co-annihilation region. Although we will not explore this question in more detail, the specific example of Table 5 shows a possibility in which the stau and neutralino are in a range of masses such that efficient co-annihilation may take place. Small variations of m_0 may lead to the precise value of the stau-neutralino mass difference necessary to obtain the proper relic density without affecting the required non-standard Higgs spectrum.

m_h [GeV]	m_A [GeV]	m_{H^+} [GeV]	$m_{\tilde{\tau}_R}$ [GeV]	$m_{\tilde{\tau}_L}$ [GeV]	$m_{\tilde{\chi}_1^0}$ [GeV]	μ [GeV]	ΔT_{tot}
200	176	255	293	1020	275	321	0.12

Table 5: Partial spectrum for $\tan \beta = 48$, $m_0 = 360$ GeV, $m_{H_u}^2 = m_{H_d}^2 = (200 \text{ GeV})^2$, $M_Y = M_2 = M_3 = 700$ GeV and $M_1 = 400$ GeV at the messenger scale.

7 Conclusions

In this article we have analyzed the phenomenological and cosmological properties of the MSSM with enhanced $SU(2)$ D-terms. Due to the splitting introduced in the third generation sfermion and non-standard Higgs boson masses, we were able to raise the Higgs mass up to about 300 GeV without being in conflict with precision electroweak tests. We worked out a particular supersymmetry breaking scenario that leads to the presence of light third generation sleptons, which provide the necessary positive contribution to the T parameter to compensate for the negative one induced by the heavy Higgs boson. Assuming universal soft supersymmetry breaking gaugino, sfermion and Higgs masses at the messenger scale, this supersymmetry breaking scenario leads to the presence of heavy squarks and non-standard Higgs bosons. The slepton spectrum is therefore determined from the requirement of reestablishing agreement with precision electroweak data.

We studied the corresponding collider signatures, which, similar to what happens in the light stau scenario in the MSSM, is characterized by the presence of many tau’s and copious missing energy in the final states. For larger values of the Higgs mass, the mass difference between the

²If very specific conditions are fulfilled, however, stau NLSP scenarios may survive and even serve to solve the so-called Lithium problem of Big Bang Nucleosynthesis [27, 28].

charged stau and the sneutrino is large enough to allow two body decays of the staus into $W^\pm$ and $\tilde{\nu}$'s. Such final states would be a signature of the enhanced $SU(2)$ D-term scenarios. Furthermore, Higgs searches into pairs of W and Z gauge bosons would be similar to the SM except in a small region of parameter space where tau sneutrinos are light enough to allow Higgs decays into sneutrino pairs. This additional decay mode may lead to a large enough modification of the branching ratios of Higgs decay into gauge bosons to avoid recent Tevatron bounds on the Higgs mass.

For the range of slepton masses consistent with precision electroweak data, the sneutrinos cannot provide a consistent thermal relic density. Indeed, mainly due to the Z -mediated annihilation channel, the thermal sneutrino dark matter density turns out to be too small. Therefore, we have assumed a gravitino LSP, heavy enough to avoid thermal equilibrium with the plasma. Gravitinos are produced by the decay of the light sneutrinos and by the scattering of particles in the early universe. For appropriate values of the reheating temperature the proper dark matter relic density may be obtained.

In the presence of a gravitino LSP, the light sneutrino decays in the early universe can lead to modifications of the light element abundances. However, for sneutrino masses leading to agreement with precision electroweak data and gravitino masses larger than a few GeV, these constraints prove to be too weak to lead to any relevant bound on the models discussed in this work.

Finally, we have discussed possible alternative scenarios in which the $SU(2)_1$ gauginos are not suppressed and therefore the third generation left-handed sleptons masses are driven to large values by their re-normalization group effects. The non-standard Higgs bosons will remain light for sufficiently large values of $\tan\beta$ and may play a relevant role in canceling the negative contributions to the T parameter induced by the heavy Higgs. This scenario is constrained by current Tevatron searches on standard and non-standard Higgs bosons and by cosmological constraints. Moreover, as is well known, the neutral and charged Higgs components may induce relevant flavor violating effects in such scenarios. However, the large mass splitting between the neutral and charged components induced by the $SU(2)_1$ D-terms leads to important differences in the flavor constraints with respect to the traditional MSSM. In particular, the heaviness of the charged Higgs dilutes the strong constraints on the MSSM proceeding from the rare decays $B^+ \rightarrow \tau^+\nu$ and $b \rightarrow s\gamma$. It would be interesting to do a more complete analysis of this question.

Acknowledgements

We would like to thank P. Batra, M. Carena, A. Delgado, E. Ponton and particularly T. Tait for useful discussions and comments. This work was supported in part by the DOE under Task TeV of contract DE-FGO3-96-ER40956. Work at UC Davis was supported in part by U.S. DOE grant No. DE-FG02-91ER40674. Work at ANL is supported in part by the US DOE, Div. of HEP, Contract DE-AC02-06CH11357. Work of N. R. Shah is supported by a Bloomenthal Research Fellowship.

References

- [1] H. P. Nilles, Phys. Rept. **110** (1984) 1; H. E. Haber and G. L. Kane, Phys. Rept. **117**(1985) 75.
- [2] S. P. Martin, arXiv:hep-ph/9709356.
- [3] H. E. Haber and R. Hempfling, Phys. Rev. Lett. **66**, 1815 (1991); Y. Okada, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. **85**, 1 (1991); J. R. Ellis, G. Ridolfi and F. Zwirner, Phys. Lett. B **262**, 477 (1991).
- [4] M. S. Carena, J. R. Espinosa, M. Quiros and C. E. M. Wagner, Phys. Lett. B **355**, 209 (1995) [arXiv:hep-ph/9504316]; M. S. Carena, M. Quiros and C. E. M. Wagner, Nucl. Phys. B **461**, 407 (1996) [arXiv:hep-ph/9508343]; H. E. Haber, R. Hempfling and A. H. Hoang, Z. Phys. C **75**, 539 (1997) [arXiv:hep-ph/9609331]; S. Heinemeyer, W. Hollik, and G. Weiglein, Phys. Rev. D **58**, 091701 (1998) [arXiv:hep-ph/9803277]; S. Heinemeyer, W. Hollik, and G. Weiglein, Phys. Lett. B **440**, 296 (1998) [arXiv:hep-ph/9807423]; S. Heinemeyer, W. Hollik, and G. Weiglein, Eur. Phys. J. C **9**, 343 (1999) [arXiv:hep-ph/9812472]; J. R. Espinosa and R. J. Zhang, J. High Energy Phys. **0003**, 026 (2000) [arXiv:hep-ph/9912236]; J. R. Espinosa and R. J. Zhang, Nucl. Phys. B **586**, 3 (2000) [arXiv:hep-ph/0003246]; M. Carena, H. E. Haber, S. Heinemeyer, W. Hollik, C. E. M. Wagner, and G. Weiglein, Nucl. Phys. B **580**, 29 (2000) [arXiv:hep-ph/0001002]; G. Degrandi, P. Slavich, and F. Zwirner, Nucl. Phys. B **611**, 403(2001) [arXiv:hep-ph/0105096]; A. Brignole, G. Degrandi, P. Slavich, and F. Zwirner, arXiv:hep-ph/0112177; S. P. Martin, Phys. Rev. D **67**, 095012 (2003) [arXiv:hep-ph/0211366].
- [5] P. Batra, A. Delgado, D. E. Kaplan and T. M. P. Tait, JHEP **0402**, 043 (2004) [arXiv:hep-ph/0309149];
- [6] P. Batra, A. Delgado, D. E. Kaplan and T. M. P. Tait, JHEP **0406**, 032 (2004) [arXiv:hep-ph/0404251].
- [7] B. Bellazzini, C. Csaki, A. Delgado and A. Weiler, arXiv:0902.0015 [hep-ph].
- [8] D. J. Muller and S. Nandi, Phys. Lett. B **383**, 345 (1996) [arXiv:hep-ph/9602390]; E. Malkawi, T. Tait and C. P. Yuan, Phys. Lett. B **385**, 304 (1996) [arXiv:hep-ph/9603349]; E. Malkawi and C. P. Yuan, Phys. Rev. D **61**, 015007 (2000) [arXiv:hep-ph/9906215].
- [9] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).
- [10] D. E. Morrissey, T. M. P. Tait and C. E. M. Wagner, Phys. Rev. D **72**, 095003 (2005) [arXiv:hep-ph/0508123].
- [11] R. Barbieri, L. J. Hall, Y. Nomura and V. S. Rychkov, Phys. Rev. D **75**, 035007 (2007) [arXiv:hep-ph/0607332].

- [12] R. Barbieri, L. J. Hall and V. S. Rychkov, Phys. Rev. D **74**, 015007 (2006) [arXiv:hep-ph/0603188].
- [13] [The ALEPH, DELPHI, L3, OPAL, SLD Collaborations, the LEP Electroweak Working Group, the SLD Electroweak and Heavy Flavour Groups], Phys. Rept. **427**, 257 (2006) [arXiv:hep-ex/0509008].
- [14] M. Takakashi [ATLAS Collaboration and CMS Collaboration], arXiv:0809.3224 [hep-ex].
- [15] J. Qian [CDF and D0 Collaborations], arXiv:0812.3979 [hep-ex].
For the most recent Tevatron Higgs search results, see
<http://theory.fnal.gov/jetp/talks/Verzocchi.pdf>
<http://theory.fnal.gov/jetp/talks/Jindariani.pdf>
- [16] A. Maloney, A. Pierce and J. G. Wacker, JHEP **0606**, 034 (2006) [arXiv:hep-ph/0409127].
- [17] J. R. Ellis, T. Falk, K. A. Olive and Y. Santoso, Nucl. Phys. B **652**, 259 (2003) [arXiv:hep-ph/0210205].
- [18] L. Covi and S. Kraml, JHEP **0708**, 015 (2007) [arXiv:hep-ph/0703130].
- [19] J. R. Ellis, K. A. Olive and Y. Santoso, JHEP **0810**, 005 (2008) [arXiv:0807.3736 [hep-ph]].
- [20] LEPSUSYWG/01-03.1 and LEPSUSYWG/02-04.1, ALEPH, DELPHI, L3 and OPAL experiments (<http://lepsusy.web.cern.ch/lepsusy/Welcome.html>).
- [21] M. Schmaltz and W. Skiba, Phys. Rev. D **62**, 095004 (2000) [arXiv:hep-ph/0004210].
- [22] M. Schmaltz and W. Skiba, Phys. Rev. D **62**, 095005 (2000) [arXiv:hep-ph/0001172].
- [23] A. D. Medina and C. E. M. Wagner, JHEP **0612**, 037 (2006) [arXiv:hep-ph/0609052].
- [24] M. Muhlleitner, A. Djouadi and Y. Mambrini, Comput. Phys. Commun. **168**, 46 (2005) [arXiv:hep-ph/0311167].
- [25] L. Cavicchia, R. Franceschini and V. S. Rychkov, Phys. Rev. D **77**, 055006 (2008) [arXiv:0710.5750 [hep-ph]].
- [26] S. P. Martin, Phys. Rev. D **75**, 115005 (2007) [arXiv:hep-ph/0703097].
- [27] R. H. Cyburt, J. R. Ellis, B. D. Fields, K. A. Olive and V. C. Spanos, JCAP **0611**, 014 (2006) [arXiv:astro-ph/0608562].
- [28] S. Bailly, K. Jedamzik and G. Moulataka, arXiv:0812.0788 [hep-ph].
- [29] K. Kadota, K. A. Olive and L. Velasco-Sevilla, arXiv:0902.2510 [hep-ph].

- [30] A. de Gouvea, T. Moroi and H. Murayama, Phys. Rev. D **56**, 1281 (1997) [arXiv:hep-ph/9701244].
- [31] T. Kanzaki, M. Kawasaki, K. Kohri and T. Moroi, Phys. Rev. D **75**, 025011 (2007) [arXiv:hep-ph/0609246].
- [32] J. L. Feng, S. Su and F. Takayama, Phys. Rev. D **70**, 075019 (2004) [arXiv:hep-ph/0404231].
- [33] M. S. Carena, M. Olechowski, S. Pokorski and C. E. M. Wagner, Nucl. Phys. B **426**, 269 (1994) [arXiv:hep-ph/9402253].
- [34] M. Carena, A. Menon and C. E. M. Wagner, arXiv:0812.3594 [hep-ph].
- [35] http://www-cdf.fnal.gov/physics/new/hdg/results/htt_070928/
<http://www-d0.fnal.gov/Run2Physics/WWW/results/prelim/HIGGS/H29/H29.pdf>